

Original Article

Incidence and Risk Factors of Embryo-logically Diagnosed Congenital Hydrocephalus: A Study of Ultrasonography Finding in 2nd and 3rd Trimesters

Riasat Ali Nahra¹, Noreen Rasul², Shabnam Sibtain³, Eisha Tahir⁴, Maham Abid⁵, Sadia Mahmood⁶

¹Professor Department of Anatomy, Pak Red Crescent Medical & Dental College, Lahore

²Associate Professor of Obs & Gynae, Pak Red Crescent Medical & Dental College, Lahore

³Assistant Professor Obs & Gynae, Azra Naheed Medical College, Lahore

^{4,5}Assistant Professor of Radiology, Shalamar Hospital, Lahore

⁶Consultant Radiologist, Lady Aitchison Hospital Lahore

Correspondence: Dr Raisat Ali Nahra

Professor Red Crescent Medical College, Lahore

drriasat@yahoo.com

Abstract

Objective: To determine the incidence of congenital hydrocephalus (CHC) detected by ultrasonography during antenatal check-ups and to assess associated risk factors in the second and third trimesters of pregnancy.

Methodology: A descriptive cross-sectional study was conducted in the Departments of Radiology and Obstetrics & Gynaecology at Pak Red Crescent Medical and Dental College and Teaching Hospital, Dina Nath, Multan Road, Kasur. Pregnant women in the second and third trimesters presenting for routine antenatal ultrasound examination, irrespective of age and parity, were included in the study. Ultrasonography was performed using high-resolution real-time ultrasound machines with colour Doppler in all participants. The ventricular atrial width was measured at the level of the glomus of the choroid plexus during scanning. Congenital hydrocephalus was considered present when the atrial width of the lateral ventricles was found to be greater than 10 mm. Data were collected using a self-structured proforma containing study-related variables.

Results: The mean maternal age was 28.34 ± 6.44 years. Congenital hydrocephalus was identified in 5 out of 2335 women, while spina bifida was detected in 3 out of 2335 women. Among the suspected risk factors, inadequate intake of folic acid, advanced maternal age, and consanguinity were found to be the most important factors and showed significant associations with both congenital hydrocephalus and spina bifida ($p < 0.05$). Hypertension, diabetes, and infections did not show any significant association with either anomaly ($p > 0.05$).

Conclusion: The incidence of congenital hydrocephalus was approximately 5 (0.2%) out of 2335 pregnancies diagnosed by ultrasonography. Inadequate folic acid intake, advanced maternal age, and consanguinity were identified as the most common suspected risk factors during the second and third trimesters of pregnancy.

Keywords: Congenital hydrocephalus, Risk factors, Ultrasonography

Cite this article as: Nahra RA, Rasul N, Sibtain S, Tahir E, Abid M, Mahmood S. Incidence and Risk Factors of Embryo-logically Diagnosed Congenital Hydrocephalus: A Study of Ultrasonography Finding in 2nd and 3rd Trimesters. J Soc Obstet Gynaecol Pak. 2026; 16(1):42-46 DOI. 0.71104/jsogp.v16i1.1033

Introduction

Congenital hydrocephalus (CH) reflects a set of multifactorial, complex, and heterogeneous disorders of brain indicated by abnormal circulation or absorption of cerebrospinal fluid (CSF), which results in ventriculomegaly (VM), a most common addressed fetal brain abnormality.¹ Although exact prevalence remains unidentified, estimates of congenital hydrocephalus cases have been suggested to range from 1% to 5% in general population, while affecting 02 to 08 neonates per 10,000 live births.^{2,3} The global prevalence of

hydrocephalus varies by geographic, genetic, and sociodemographic factors.²⁻⁴ Global annual incidence of hydrocephalus among infants has been estimated to be over 383,000 fresh cases, with case-fatality levels reflecting 4%-87% deaths among individuals affected by hydrocephalus.⁴ A range of etiological factors can contribute to congenital hydrocephalus development, including vitamin deficiency, viral infections, developmental abnormalities, genetic predisposition, and environmental factors.^{5,6} In Pakistan around 6% to

Authorship Contribution:^{1,5} Substantial contributions to the conception or design of the work or the acquisition, ²Final approval of the study to be published, ^{4,6}Drafting the work or revising it critically for important intellectual content.

Funding Source: none

Conflict of Interest: none

Received: July 18, 2025

Revised: Nov 05, 2025

Accepted: Dec 11, 2025

9% of perinatal deaths are attributed to congenital conditions and these anomalies have been associated with maternal, genetic, sociodemographic factors, and metabolic and cardiovascular conditions.⁷

Ultrasonography remains a major diagnostic modality for the detection of cerebral ventriculomegaly and congenital hydrocephalus in obstetrical practice due to its convenience, safety, and acceptable diagnostic accuracy, particularly during the second and third trimesters compared with the first trimester.⁸⁻¹⁰ In this regard, standardized measurement of the lateral ventricles for the early identification of ventriculomegaly facilitates monitoring of disease progression, detection of associated anomalies, and assists in clinical decision-making regarding prognosis and management.¹¹

A recent study reported the detection of various brain anomalies on prenatal ultrasonography, including neural tube defects in 75.3% of cases, isolated central nervous system anomalies in 74.3%, combined central nervous system and extracranial anomalies in 5.2% and 21.9%, respectively, and ventriculomegaly in 11.0% of cases, with an overall mortality rate reaching up to 43%.¹²

Despite advancements in prenatal imaging techniques and diagnostic approaches, findings across studies remain heterogeneous. This highlights the need for population-based research incorporating second- and third-trimester sonographic evaluations to determine more accurate estimates of the incidence of congenital hydrocephalus and its associated risk factors. Therefore, the present study was conducted to determine the incidence of congenital hydrocephalus and to identify related maternal and fetal risk factors in our population. Such evidence may help improve early diagnosis, support the development of preventive strategies, and enhance antenatal counselling and management, ultimately contributing to better fetomaternal outcomes.

Methodology

This descriptive cross-sectional study was conducted in the Departments of Radiology and Obstetrics & Gynaecology at Pak Red Crescent Medical College and Teaching Hospital, Lahore. Study duration was one year from July 2023 to June 2024. All pregnant women in the second and third trimesters with singleton pregnancy, presenting for routine antenatal ultrasound examination, irrespective of age and parity, were included in the study. Women with multiple gestations (twin pregnancies), poor visualization due to maternal obesity or fetal position, the

presence of other major fetal anomalies, and those who were unwilling to participate were excluded.

A non-probability consecutive sampling technique was employed, and all eligible participants fulfilling the inclusion criteria were enrolled until the required sample size was achieved. Ultrasonographic examinations were performed using high-resolution real-time ultrasound machines with colour Doppler by a well-trained and experienced sonologist with a minimum of five years of professional experience. Standard obstetric ultrasound scanning protocols were followed. Particular attention was given to the evaluation of the third ventricle, lateral ventricles, cavum septi pellucidi, cerebral hemispheres, and posterior fossa structures.

The ventricular atrial width was measured at the level of the glomus of the choroid plexus during scanning. Congenital hydrocephalus was considered present when the atrial width of the lateral ventricles was found to be greater than 10 mm, either with or without associated findings such as a dangling choroid plexus, thinning of the cortical mantle, or enlargement of the third ventricle. Obstetric risk factors were assessed through direct maternal history and review of antenatal medical records. Data were collected using a self-structured proforma containing study-related variables, and statistical analysis was performed using SPSS version 23.

Results

The maternal average age was 28.34 ± 6.44 years, and average gestational age was 24.31 ± 5.29 weeks. Around More than half of the women were presented during the 2nd trimester (54.2%), while 45.8% were presented during third trimester of pregnancy. Out of all, 28.7% of cases had consanguinity; the satisfactory use of folic acid was noted 61.0% of women during pregnancy, while rest had history of inadequately intake of folic acid. The antenatal infections were found in 13.2% of the women, whereas the hypertensive and diabetics women were 19.2% and 14.3%, respectively. (Table I)

Out of one year presented pregnancies (2335) were evaluated by ultrasonography, where congenital hydrocephalus was found around 5/2335 (0.2%) of the women, while spina bifida was detected among 3 out of 2335 (0.1%) of the women. (Table II)

According to the suspected risk factors the inadequately given folic acid and advance maternal age were the found most important risk factors and which showed

significant correlations with both congenital hydrocephalus and the spina bifida $p < 0.05$. The consanguinity was also significantly associated with

Table I: Demographic and clinical variables of the patients. (n=2335)

Variables	N	%
Consanguinity	No	1664
	Yes	671
Use of folic acid	Adequate	1424
	Inadequate	911
Antenatal infection	No	2026
	Yes	309
Diabetes mellitus	No	2002
	Yes	333
Hypertension	No	1886
	Yes	449
Trimester of pregnancy	Second	1265
	Third	1070
Maternal mean age	28.34±6.44 years	
Mean gestational age	24.31±5.29 weeks	

Table II: Frequency of congenital disorders. (n=2335)

Congenital disorders	N	%
Congenital Hydrocephalus		
No	2330	99.8
Yes	05	0.2
Total	2335	100.0
Spina Bifida		
No	2332	99.9
Yes	03	0.1
Total	2335	100.0

congenital hydrocephalus only ($p=0.011$). On the other hand, diabetes, and infections did not show the any significant correlation with either anomalies, $p > 0.05$ ($p > 0.05$), indicating the consanguinity, advanced maternal age and poor periconceptional folic acid intake play a key role congenital anomalies development. (Table III)

Discussion

This study included pregnant women in the second and third trimesters. The mean maternal age was 28.34 ± 6.44 years, and the mean gestational age was 24.31 ± 5.29 weeks. More than half of the women presented during the second trimester (54.7%), whereas 45.8% presented during the third trimester of pregnancy. Comparable findings were reported in a study conducted by Alia and Ahmed¹³, who evaluated antenatal ultrasound examinations in the second and third trimesters and documented a mean maternal age of 26.5 years and a mean gestational age of 24 weeks at diagnosis. Similarly, another study by Mahela and Talukdar¹⁴ reported a mean maternal age of 25.5 ± 6.15 years and a mean gestational age of 27 ± 6.42 weeks, suggesting a relatively late presentation of pregnant women for routine antenatal screening.

Table III: Analysis for post stratification. (n=2335)

Variables		Congenital Hydrocephalus		Total	p-value	Spina Bifida		Total	p-value
		No	Yes			No	Yes		
Age groups	18-25 years	1013	1	1014	0.043	1014	0	1014	0.028
		43.4%	0.0%	43.4%		43.4%	0.0%	43.4%	
	26-32 years	629	0	629		629	0	629	
		26.9%	0.0%	26.9%		26.9%	0.0%	26.9%	
	33-40 years	688	4	692		689	3	692	
		29.5%	0.2%	29.6%		29.5%	0.1%	29.6%	
Consanguinity	No	1663	1	1664	0.011	1663	1	1664	0.146
		71.2%	0.0%	71.3%		71.2%	0.0%	71.3%	
	Yes	669	2	671		669	2	671	
		28.7%	0.1%	28.7%		28.7%	0.1%	28.7%	
Diabetes mellitus	No	1998	4	2002	0.713	1999	3	2002	0.480
		85.6%	0.2%	85.7%		85.6%	0.1%	85.7%	
	Yes	332	1	333		333	0	333	
		14.2%	0.0%	14.3%		14.3%	0.0%	14.3%	
Hypertension	No	1883	3	1886	0.238	1885	1	1886	0.037
		80.6%	0.1%	80.8%		80.7%	0.0%	80.8%	
	Yes	447	2	449		447	2	449	
		19.1%	0.1%	19.2%		19.1%	0.1%	19.2%	
Folic acid use	adequate	1424	0	1424	0.005	1424	0	1424	0.030
		61.0%	0.0%	61.0%		61.0%	0.0%	61.0%	
	Inadequate	906	5	911		908	3	911	
		38.8%	0.2%	39.0%		38.9%	0.1%	39.0%	
Antenatal infection	No	2024	2	2026	0.062	2025	1	2026	0.076
		86.7%	0.1%	86.8%		86.7%	0.0%	86.8%	
	Yes	306	3	309		307	2	309	
		13.1%	0.1%	13.2%		13.1%	0.1%	13.2%	

According to the current study, out of all, 28.7% of cases had consanguinity, the satisfactory use of folic acid was noted 61.0% of women during pregnancy, while rest had history of inadequately intake of folic acid. Aligning with these findings, in the study of Khan et al.,¹⁵ who studies congenital anomalies among infants and reported consanguinity rate of 67.7%, which is higher than that of our study, and folic acid supplementation in 32.3% of women during pregnancy, which is lower than our finds. Similarly, in the study of Karim et al.¹⁶ higher consanguinity rate of 83.9% was found than our study findings, with 77.5% of women never consumed folic acid supplements during pregnancy, reflecting common pattern of lack of folic acid use. In current study, the antenatal infections were found in 13.2% of the women, whereas the hypertensive and diabetics women were 19.2% and 14.3%, respectively. Consistent findings were observed in the study carried out by Nahar et al.,¹⁷ who found gestational/uncontrolled diabetes among 16% and additional maternal conditions such as representing rash accompanied by fever in 12% of patients. Consistently in the study of Anbreen et al.,¹⁸ diabetes was present in 5.57% of patient, which was relatively lower than our findings.

In present study, out of one year presented pregnancies were evaluated by ultrasonography, where congenital hydrocephalus was found in around 0.2% of the women, while spina bifida was detected among 0.9% of the women. In agreement with these findings, in the study conducted by Mahela and Talukdar¹⁹ congenital abnormalities were found in 1.57% of pregnant women in their second or third trimester. Aligning with these findings, in a study conducted by Mahmoud et al.²⁰ hydrocephalus was detected in 0.4% of cases and spina bifida in 30% of hydrocephalus cases. Incidence of hydrocephalus their study was lower, while spina bifida cases were higher than those of our study. In our study cohort, according to the suspected risk factors the inadequately given folic acid and consanguinity were the found most important risk factors and which showed significant correlations with both congenital hydrocephalus and the spina bifida $p < 0.05$. Aligning with our study findings, in the study carried out by Maged et al.²¹ supplementation of folate was significantly associated with spina bifida ($p < 0.001$). They investigated consanguinity as a risk factor and did not find statistical significance. In another study conducted by Saleem et al.²² periconceptional supplementation of folic acid and high parental consanguinity were found to be significantly associated with neural tube defect ($p <$

0.01), suggesting that consanguinity and inadequate folic acid use are important risk factors in congenital anomalies.

In this study, the age of mothers, hypertension, diabetes, trimester of pregnancy and infections did not show the any significant correlation with either anomalies, $p > 0.05$ ($p > 0.05$), indicating the consanguinity and poor periconceptional folic acid intake play a key role congenital anomalies development. In agreement with our findings, in the study of Karim et al.²³ although, inadequate use of folic and consanguinity were the most frequent risk factors, use of folic, consanguinity, maternal age and diabetes were not significantly correlated with congenital anomalies ($p > 0.05$).

Conclusion

Incidence of CHC was observed around 5(0.2%)/2335 pregnancy diagnosed by ultrasonography during 2nd and 3rd trimester of pregnancy, along with the significance risk factors as low folic acid intake, advanced maternal age and three cases of spina bifida were found with neural tube defect also related to folic acid deficiency.

References

1. Siddiqui RB, Erum U, Rauf R, Shakoar S, Zulfiqar S, Razzaq M. Maternal environmental factors associated with congenital hydrocephalus. *J Fatima Jinnah Med Univ.* 2024;18(1):25-8. <https://doi.org/10.37018/JFJMU/ADN/6524>
2. Shakeel NA, Aziz A, Farooq MF, Usmani F, Salahuddin T. Obstructive hydrocephalus: short term complications of endoscopic third ventriculostomy in obstructive hydrocephalus. *Prof Med J.* 2019;26(2):318-23. <https://doi.org/10.29309/TPMJ/2019.26.02.3115>
3. Jacquemin V, Versbaegen N, Duerinckx S, Massart A, Soblet J, Perazzolo C, et al. Congenital hydrocephalus: new Mendelian mutations and evidence for oligogenic inheritance. *Hum Genomics.* 2023;17(1):16. doi: <https://doi.org/10.1186/s40246-023-00464-w>
4. Dewan MC, Rattani A, Mekary R, Glancz LJ, Yunusa I, Baticulon RE, et al. Global hydrocephalus epidemiology and incidence: systematic review and meta-analysis. *J Neurosurg.* 2019;130(4):1065-79. <https://doi.org/10.3171/2017.10.JNS17439>
5. Liu XY, Song X, Czosnyka M, Robba C, Czosnyka Z, Summers JL, et al. Congenital hydrocephalus: a review of recent advances in genetic etiology and molecular mechanisms. *Mil Med Res.* 2024;11(1):54. doi: <https://doi.org/10.1186/s40779-024-00560-5>
6. Karim JK, Reeves BC, Damisah E, Duy PQ, Antwi P, David W, et al. Inflammation in acquired hydrocephalus: pathogenic mechanisms and therapeutic targets. *Nat Rev Neurol.* 2020;16(5):285-96. <https://doi.org/10.1038/s41582-020-0321-y>
7. Awan JA, Qamar A, Surti A, Anwar E. Are congenital malformations associated with maternal sociodemographic and risk factors? A multicenter ultrasound-based study. *Pak J Med Sci.* 2024;40(11):2538. doi: <https://doi.org/10.12669/pjms.40.11.9792>
8. Feroz S, Ali N, Hasan ZU, Fatima Z, Hina N, Jahangir A, et al. Ultrasonographic findings of ventriculomegaly in second and third trimester with fetal outcomes. *Pak J Health Sci.* 2023;4(2):27-31. <https://doi.org/10.54393/pjhs.v4i02.465>

9. Routh ML. Role of ultrasound and its recent applications in diagnosing fetal malformations. In: *Selected Topics in Prenatal and Neonatal Diagnoses*. IntechOpen; 2024.
10. Buijtendijk MF, Bet BB, Leeftang MM, Shah H, Reuvekamp T, Goring T, et al. Diagnostic accuracy of ultrasound screening for fetal structural abnormalities during the first and second trimester of pregnancy in low-risk and unselected populations. *Cochrane Database Syst Rev*. 2024;(5). doi: <https://doi.org/10.1002/14651858.CD014715.pub2>
11. D'Addario V, Rossi A. Mild fetal ventriculomegaly: diagnostic work-up and management. *Donald Sch J Ultrasound Obstet Gynecol*. 2011.
12. Bağcı M, Karaaslan O, Başkiran Y. Evaluation of fetal central nervous system anomalies diagnosed prenatally: prenatal and postnatal outcomes. *Van Tip Derg*. 2024;31(2). <https://doi.org/10.5505/vmj.2024.77487>
13. Alia N, Ahmed I. Congenital anomalies: prevalence of congenital abnormalities in second trimester of pregnancy in Madina Teaching Hospital Faisalabad on gray scale ultrasound. *J Univ Med Dent Coll*. 2010;1(1):23-8.
14. Mahela S, Talukdar B. Prevalence of congenital abnormalities on routine ultrasound scan of second and third trimester pregnancy. *Int J Reprod Contracept Obstet Gynecol*. 2016;5(1):182-5. <https://doi.org/10.18203/2320-1770.ijrcog20151621>
15. Khan A, Zuhaid M, Fayaz M, Ali F, Khan A, Ullah R, et al. Frequency of congenital anomalies in newborns and its relation to maternal health in a tertiary care hospital in Peshawar, Pakistan. *Int J Med Stud*. 2015;3(1):19-23. doi: <https://doi.org/10.5195/ijms.2015.108>
16. Karim R, Wahab S, Akhtar R, Jamala F, Jabeen S. Frequency and pattern of distribution of antenatally diagnosed congenital anomalies and associated risk factors. *J Postgrad Med Inst*. 2014;28(2).
17. Nahar N, Ony FK, Liza TR, Jahan N, Bishnu S. Risk factors associated with fetal congenital anomaly among pregnant women. *TAJ J Teach Assoc*. 2024;37(2):724-30. <https://doi.org/10.70818/taj.v037i02.0507>
18. Anbreen T, Ali L, Butt S, Shah T. Congenital anomaly frequency, risk factor and trends among antenatal patients presenting at tertiary care hospital in Pakistan. *Pak J Med Res*. 2021;60(2):52-6.
19. Mahela S, Talukdar B. Prevalence of congenital abnormalities on routine ultrasound scan of second and third trimester pregnancy. *Int J Reprod Contracept Obstet Gynecol*. 2016;5(1):182-5. <https://doi.org/10.18203/2320-1770.ijrcog20151621>
20. Mahmoud MZ, Dinar HA, Abdulla AA, Babikir E, Sulieman A. Study of the association between the incidences of congenital anomalies and hydrocephalus in Sudanese fetuses. *Glob J Health Sci*. 2014;6(5):1. <https://doi.org/10.5539/gjhs.v6n5p1>
21. Maged A, Elsherbini M, Ramadan W, Elkomy R, Helal O, Hatem D, et al. Periconceptional risk factors of spina bifida among Egyptian population: a case-control study. *J Matern Fetal Neonatal Med*. 2016;29(14):2264-7. <https://doi.org/10.3109/14767058.2015.1081890>
22. Saleem MU, Ali MU, Hussnain QA. Prevalence and associated risk factors for neural tube defects in patients at Sheikh Zayed Hospital Rahim Yar Khan. *Pak J Med Sci*. 2010;4:197-202.
23. Karim R, Wahab S, Akhtar R, Jamala F, Jabeen S. Frequency and pattern of distribution of antenatally diagnosed congenital anomalies and associated risk factors. *J Postgrad Med Inst*. 2014;28(2).